

Tele-Medicine for Cardiovascular Emergencies: Ready for the Prime Time?

Natale Daniele Brunetti^{1*}, Daniele Amoroso², Luisa De Gennaro¹, Giulia Dellegrottaglie³, Giuseppe Di Giuseppe³, Gianfranco Antonelli⁴ and Matteo Di Biase⁵

¹Post-graduate in Cardiology, PhD, Cardiology Department, University of Foggia, Italy

²Post-graduate in Radiology, U.O. Cardiologia, Azienda Ospedaliera Policlinico, Bari, Italy

³Cardio on Line Europe S.R.L., Bari, Italy

⁴Post-graduate in Cardiology, Cardiology Department, Azienda Ospedaliera Policlinico, Bari, Italy

⁵Post-graduate in Cardiology, Cardiology Department, University of Foggia, Italy

Abstract

According to the Wikipedia definition, "Tele-Medicine [(TM)] is the use of telecommunication and information technologies in order to provide clinical health care at a distance. It helps eliminate distance barriers and can improve access to medical services that would often not be consistently available in distant rural communities. It is also used to save lives in critical care and emergency situations". TM may therefore be extremely useful in reducing time of treatment, especially when time is paramount, such as in cardiovascular emergencies.

Pre-hospital Electrocardiograms (ECGs) may be useful in improving timeliness of treatments in acute cardiovascular disease, and TM may play a primary role in allowing any patient with a suspected cardiovascular emergency to be examined with pre-hospital ECG by a cardiologist, wherever the patient is, either in rural or urban areas. In the field of TM applied to cardiovascular emergencies managed by an Emergency Medical Service (EMS) in a large region-wide area, the focus is on Apulia, a region in South-Eastern Italy (19,358 km², population 4,091,259). The region is currently served by a TM *hub* from where cardiologists report back, 24/7, on pre-hospital ECGs transmitted by telephone (mobile or other). Evidence is provided showing the clinical utility of pre-hospital TM ECG in improving the quality of medical assistance in cardiovascular emergencies.

The Apulia TM network provides an interesting demonstration of TM potential in health-care management, combining high-quality medical assistance with optimal spending of public resources.

Keywords: Tele-Medicine; Cardiology; Emergency medical service

According to the Wikipedia definition, "tele-medicine (TM) is the use of telecommunication and information technologies in order to provide clinical health care at a distance. It helps eliminate distance barriers and can improve access to medical services that would often not be consistently available in distant rural communities. It is also used to save lives in critical care and emergency situations" [1]. So TM may prove extremely useful in overcoming the problem of distance in emergency situations and then reducing delay to treatment when time is paramount, and time is paramount especially in cardiovascular emergencies.

There is evidence that, when treating patients with Acute Myocardial Infarction (AMI), the time from the onset of symptoms to reperfusion is crucial for salvaging myocardium; in other words, "time is muscle". Shorter time-to-reperfusion is associated with smaller infarct size and micro-vascular damage and larger salvaged myocardium [2]. Fast reperfusion may also reduce both in-hospital rates of adverse events after AMI [3], and during the following 12 months [4]. A shorter symptom-to-balloon time is associated with an improved coronary flow, an increased likelihood of subsequent left ventricular systolic ejection fraction > 40 %, and greater 3-year survival [5]. The higher the risk profile, the larger the reduction in mortality benefits from fast reperfusion [6].

However, of 4278 patients transferred for primary angioplasty (PCI) in the United States, only 4-10% of patients were treated within 90 minutes, the benchmark recommended by national quality guidelines [7,8].

Several strategies are currently suggested for achieving better adherence to international guidelines on management of AMI and

timely myocardial reperfusion. In the Vienna registry, an example of a good hospital network with prompt PCI or fibrinolysis availability, a marked increase in the proportion of patients receiving PCI, and a reduction in the proportion receiving no reperfusion therapy, was achieved [9].

Pre-hospital electrocardiograms (ECG) may be useful in decreasing delay to treatment in patients with AMI [10-14]. International guidelines, consensus and scientific statements [15] recommend that emergency medical services (EMS) acquire and use pre-hospital ECGs, and many communities are thus implementing pre-hospital ECG programs.

TM may play a starring role in allowing all patients with suspected cardiovascular emergencies to be examined with pre-hospital ECG by a cardiologist, wherever those patients are, either in rural or urban areas.

In the field of TM applied to cardiovascular emergencies and EMS in large region-wide areas, the focus is on Apulia, a region in

***Corresponding author:** Natale Daniele Brunetti, MD, post-graduate in Cardiology, PhD, Cardiology Department, University of Foggia, Viale Pinto 1, 71100 Foggia, Italy, Tel: +393389112358; Fax: +390881745424; E-mail: nd.brunetti@unifg.it

Received June 07, 2012; **Accepted** July 02, 2012; **Published** July 05, 2012

Citation: Brunetti ND, Amoroso D, Gennaro LD, Dellegrottaglie G, Giuseppe GD, et al. (2012) Tele-Medicine for Cardiovascular Emergencies: Ready for the Prime Time? J Clin Exp Cardiol 3:e109. doi:[10.4172/2155-9880.1000e109](https://doi.org/10.4172/2155-9880.1000e109)

Copyright: © 2012 Brunetti ND, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

South-Eastern Italy (19,358 km², population 4,091,259). The region is currently served by a TM *hub* from where cardiologists, available 24/7, report back on ECGs recorded on a small pocket device, the CardioVox P12™ (Aerotel, Holon, Israel), and transmitted by telephone (mobile or other). The device may record a complete 12-lead ECG in a couple of minutes; after telephone coupling, the ECG is sent to the TM *hub*, which is located in Puglia's capital city, Bari. Here, a cardiologist may read the ECG on a computer screen and report back on the findings; when necessary, a brief consultation may also occur between the on-site sender – a physician, paramedics or others – and the cardiologist. Remote senders may receive back the ECG report on their smart-phone or iPad, or on a PC via an internet or telephone connection. The ECGs may be easily stored and printed as required. The TM service has been in operation since 2004 and is available for the “118” regional public EMS, the local “911”, and hundreds of private customers (patients, pharmacies, general practitioners, specialists in occupational medicine, nursing homes).

In June 2012 the 500,000th ECG will be analysed by the service, which is a particularly interesting example of synergy between public health authorities and private enterprise. The regional 118 service is run by the regional government while the tele-cardiology service is provided by Cardio-on-line Europe s.r.l., a private company.

The tele-cardiology service in Apulia combines two principal models of TM delivery for cardiology emergencies: the Boston model, with few TM providers, small areas and paramedic-based ECG analysis, and the Los Angeles model, with a larger area but a large number of TM providers and a computer-algorithm-based ECG report [16]. The Apulia model provides not only qualified, remote, on-site assistance from a cardiologist, as is required by Italian law for ECG analysis, but also cost reduction: a single TM *hub* may potentially support even more than one region with several million inhabitants.

Several studies have shown that TM support can improve the quality of medical assistance and reduce cardiovascular mortality. Transmission of a pre-hospital 12-lead ECG directly to the attending cardiologist's mobile telephone decreased door-to-PCI time by >1 hour when patients were transported directly to PCI centres, bypassing local hospitals.

At a local level, in Apulia, over 233,657 pre-hospital ECGs in subjects with suspected acute heart disease screened using pre-hospital ECG through TM support, 45% were abnormal; of these ECGs, 18% showed signs suggestive of Acute Coronary Syndrome (ACS), 20% were indicative of arrhythmia, and minor problems emerged in 62% of cases [17]. In cases of suspected ACS, ECG findings were normal in 77% of patients; 74% of subjects with suspected ACS were screened within 30' from the onset of symptoms.

In another study, ECG showed signs of ST-elevation in only 3.84% of patients with acute chest pain, while only 78.94% of patients with STEMI reported acute chest pain. The diagnosis was therefore based mainly on ECG findings with TM pre-hospital ECGs despite the presence of atypical clinical presentation [18]. About two thirds of patients with STEMI were resident in small towns without coronary care units, and thus particularly benefitted from an immediate pre-hospital diagnosis.

TM pre-hospital ECG screening could significantly help in avoiding errors and delay in STEMI diagnosis in elderly patients [19]. Among STEMI patients older than 70 years, atypical presentation was detected in 32% of patients (vs. 11%). Number needed to diagnose a STEMI with atypical clinical presentation was 3 among subjects older than 70 years.

TM support may also improve the sensitivity of diagnosis of atrial fibrillation with atypical presentation in elderly patients [20], up to 4-to-7 fold.

So tele-cardiology is now well established in Apulia. Several studies of primary and secondary prevention, also with the support of pharmacies, are currently under way. In the CAPITAL (Cardiovascular Prevention with Telecardiology in Apulia) study, 10 thousand subjects are currently involved in a project aimed at estimating cardiovascular risk and charting cardiovascular risk in a Mediterranean context [21].

Screening studies for early ECG diagnosis of cardiovascular disease among primary and high school students and professional athletes are currently in the advanced planning phase. Further initiatives will probably be useful in the future in reducing waiting lists for ECG examinations in public regional hospitals.

The Apulia TM network thus provides an interesting demonstration of TM potential in health care management, combining high-quality medical assistance with optimal spending of public resources. Tele-cardiology is ready for the prime time in cardiovascular emergencies.

References

1. <http://en.wikipedia.org/wiki/Telemedicine>
2. Francone M, Bucciarelli-Ducci C, Carbone I, Canali E, Scardala R, et al. (2009) Impact of primary coronary angioplasty delay on myocardial salvage, infarct size, and microvascular damage in patients with ST-segment elevation myocardial infarction: insight from cardiovascular magnetic resonance. *J Am Coll Cardiol* 54: 2145-2153.
3. Nielsen PH, Terkelsen CJ, Nielsen TT, Thuesen L, Krusell LR, et al. (2011) System delay and timing of intervention in acute myocardial infarction (from the Danish Acute Myocardial Infarction-2 [DANAMI-2] trial). *Am J Cardiol* 108: 776-781.
4. Fu Y, Goodman S, Chang WC, Van De Werf F, Granger CB, et al. (2001) Time to treatment influences the impact of ST-segment resolution on one-year prognosis: insights from the assessment of the safety and efficacy of a new thrombolytic (ASSENT-2) trial. *Circulation* 104: 2653-2659.
5. Maeng M, Nielsen PH, Busk M, Mortensen LS, Kristensen SD, et al. (2010) Time to treatment and three-year mortality after primary percutaneous coronary intervention for ST-segment elevation myocardial infarction—a DANish Trial in Acute Myocardial Infarction-2 (DANAMI-2) substudy. *Am J Cardiol* 105: 1528-1534.
6. De Luca G, Casseti E, Marino P (2009) Percutaneous coronary intervention-related time delay, patient's risk profile, and survival benefits of primary angioplasty vs lytic therapy in ST-segment elevation myocardial infarction. *Am J Em Med* 27: 712-719.
7. Nallamothu BK, Bates ER, Herrin J, Wang Y, Bradley EH, et al. (2005) Times to treatment in transfer patients undergoing primary percutaneous coronary intervention in the United States: National Registry of Myocardial Infarction (NRMII)-3/4 analysis. *Circulation* 111: 761-767.
8. Wang TY, Peterson ED, Ou FS, Nallamothu BK, Rumsfeld JS, et al. (2011) Door-to-balloon times for patients with ST-segment elevation myocardial infarction requiring interhospital transfer for primary percutaneous coronary intervention: a report from the national cardiovascular data registry. *Am Heart J* 161: 76-83.
9. Kalla K, Christ G, Karnik R, Malzer R, Norman G, et al. (2006) Vienna STEMI Registry Group Implementation of guidelines improves the standard of care: the Viennese Registry on Reperfusion Strategies in ST-Elevation Myocardial Infarction. *Circulation* 113: 2398-2405.
10. Morrow DA, Antman EM, Sayah A, Schuhwerk KC, Giugliano RP, et al. (2002) Evaluation of the time saved by prehospital initiation of reteplase for ST-elevation myocardial infarction: results of the Early Reteplase-Thrombolysis in Myocardial Infarction (ER-TIMI) 19 trial. *J Am Coll Cardiol* 40: 71-77.
11. Ferguson JD, Brady WJ, Perron AD, Kielar ND, Benner JP, et al. (2003) The prehospital 12-lead electrocardiogram: impact on management of the out-of-hospital acute coronary syndrome patient. *Am J Emerg Med* 21: 136-142.

12. Terkelsen CJ, Lassen JF, Nørgaard BL, Gerdes JC, Poulsen SH, et al. (2005) Reduction of treatment delay in patients with ST-elevation myocardial infarction: Impact of pre-hospital diagnosis and direct referral to primary percutaneous coronary intervention. *Eur Heart J* 26: 770–777.
13. Brainard AH, Raynovich W, Tandberg D, Bedrick EJ (2005) The prehospital 12-lead electrocardiogram's effect on time to initiation of reperfusion therapy: a systematic review and meta-analysis of existing literature. *Am J Emerg Med* 23: 351–356.
14. Curtis JP, Portnay EL, Wang Y, McNamara RL, Herrin J, et al. (2006) The pre-hospital electrocardiogram and time to reperfusion in patients with acute myocardial infarction, 2000–2002: findings from the National Registry of Myocardial Infarction-4. *J Am Coll Cardiol* 47: 1544–1552.
15. Ting HH, Krumholz HM, Bradley EH, Cone DC, Curtis JP, et al. (2008) Implementation and integration of prehospital ECGs into systems of care for acute coronary syndrome: a scientific statement from the American Heart Association Interdisciplinary Council on Quality of Care and Outcomes Research, Emergency Cardiovascular Care Committee, Council on Cardiovascular Nursing, and Council on Clinical Cardiology. *Circulation* 118: 1066–1079.
16. Sejersten M, Sillesen M, Hansen PR, Nielsen SL, Nielsen H, et al. (2008) Effect on treatment delay of prehospital teletransmission of 12-lead electrocardiogram to a cardiologist for immediate triage and direct referral of patients with ST-segment elevation acute myocardial infarction to primary percutaneous coronary intervention. *Am J Cardiol* 101: 941-946.
17. Brunetti ND, De Gennaro L, Dellegrottaglie G, Amoroso D, Antonelli G, et al. (2011) A regional prehospital electrocardiogram network with a single telecardiology “hub” for public emergency medical service: technical requirements, logistics, manpower, and preliminary results. *Telemed J E Health* 17: 727-733.
18. Brunetti ND, Amodio G, De Gennaro L, Dellegrottaglie G, Pellegrino PL, et al. (2009) Telecardiology applied to a region-wide public emergency health-care service. *J Thromb Thrombolysis* 28: 23-30.
19. Brunetti ND, De Gennaro L, Amodio G, Dellegrottaglie G, Pellegrino PL, et al. (2010) Telecardiology improves quality of diagnosis and reduces delay to treatment in elderly patients with acute myocardial infarction and atypical presentation. *Eur J Cardiovasc Prev Rehabil* 17: 615-620.
20. Brunetti ND, De Gennaro L, Pellegrino PL, Dellegrottaglie G, Antonelli G, et al. (2012) Atrial fibrillation with symptoms other than palpitations: incremental diagnostic sensitivity with at-home tele-cardiology assessment for emergency medical service. *Eur J Prev Cardiol* 19: 306-313.
21. Brunetti ND, De Gennaro L, Dellegrottaglie G, Straziota E, Novielli V, et al. (2011) Rationale and design for a cardiovascular screening and prevention study with tele-cardiology in Mediterranean Italy: the CAPITAL study (CArdiovascular Prevention with Telecardiology in ApuLia). *Int J Cardiol* 149: 130-133.

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 200 Open Access Journals
- 15,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.editorialmanager.com/clinicalgroup